

# Perilous Puddles

Admittedly, the research subject isn't particularly appetizing: *Strongyloides stercoralis* – small parasitic worms that live in their host's intestines and have the potential to cause severe problems. Nevertheless, **Adrian Streit** from the **Max Planck Institute for Developmental Biology** in Tübingen is fascinated by this threadworm. It has a unique life cycle, and to this day, no one really understands why.

TEXT **CATARINA PIETSCHMANN**

**R**eferring to nematodes as unusual is almost an understatement, as strange behavior is completely normal for them. *Pristionchus pacificus*, which lives, among other places, on the Pacific island of La Réunion, seeks out a beetle larva, climbs up onto it and then stops developing. As soon as the beetle dies, the worm continues its development, gorges itself on the carcass and multiplies (MAXPLANCKRESEARCH 2/2014).

But compared with *Strongyloides*, that's almost boringly conventional. At the Max Planck Institute for Developmental Biology, Adrian Streit explores how this worm can survive in two worlds. Between parasitic generations, *Strongyloides* can also form free-living generations. Parasites are exclusively females that multiply in the host's intes-

tines by parthenogenesis. "They produce both male and female eggs, which are excreted with the feces," explains Streit. "Either infectious larvae develop from the female eggs, which then immediately crawl back into the host, or free-living worms that mate with males are produced."

## PARASITIC FEMALES

The male worms are exclusively free-living. If males and females reproduce in the soil, only parasitic female offspring are formed. This second generation of larvae thus needs to find another host to be able to multiply – which then once again takes place without a male partner.

Many terrestrial vertebrate species have their own *Strongyloides* – including humans. The World Health Organiza-

tion estimates that more than 300 million people worldwide are infected with the threadworm, especially in northern South America, Central Africa and Asia. A warm, humid climate and a lack of hygiene are an El Dorado for worms!

In healthy people, the infection usually goes undetected, as they harbor only comparatively few worms. In immunocompromised patients, however, the larvae in the intestines can infect other organs and cause a life-threatening infection called strongyloidiasis. "If the parasite isn't detected in cancer patients, for example, chemotherapy can end in disaster," emphasizes Streit.


Organ recipients are also at risk. Even in Western countries, deaths have occurred after transplants as a result of the worms. In the Netherlands, two cases were reported in which the infection was proven to have been transmitted





Walking barefoot where the soil is contaminated with animal and human excrement should be avoided at all costs. Some parasites, such as the threadworm *Strongyloides stercoralis*, bore through their hosts' skin. The worm reproduces in the intestines and returns to the environment via the feces.





*Strongyloides stercoralis* life cycle: In a host, all worms are female. They produce female and male progeny by parthenogenesis (L1 to L4: larval stages 1 to 4). Females can enter one of three cycles: direct, indirect or auto-infectious. In the latter, the host is reinfected by parasites already present in the body. Males, in contrast, live exclusively in the indirect cycle and are free-living. All progeny from the indirect cycle are female and become parasites.

via donated organs – the donor had lived in South America 20 years previously. “In this country, the worms are not yet a major medical problem, but these cases are reason enough to take a closer look,” says Streit.

It isn’t unusual for the infection to remain undetected for a long time, as it can be completely symptom-free. In addition, the classic symptoms – skin rash, nausea, diarrhea, abdominal cramps – are unspecific, so it’s easy to overlook the worms. The tragedy is that common worm remedies would have been sufficient to kill the parasites.

A lack of sanitary hygiene results in the worm being transmitted from person to person. But is this really the only way? Adrian Streit is driven by the question of whether strongyloidiasis is one of the zoonoses, meaning that it can be transmitted from ani-

mals – dogs, for example – to humans. If this is the case, worms with identical DNA would have to be found in both dogs and their owners.

### FIELD STUDY IN CAMBODIA

To learn more about the transmission paths, Streit has teamed up with the Swiss Tropical and Public Health Institute in Basel and the Cambodian National Center for Parasitology, Entomology and Malaria Control. The two institutes maintain a field laboratory in northern Cambodia. The rural region is ideal for this purpose: the farmers’ houses are built on stilts – the family lives upstairs, and the animals, generally pigs and dogs, live below. The sanitary facilities are anything but hygienic and residents go barefoot or wear open sandals.

Nematodes survive for weeks in the moist, feces-contaminated soil. The larvae bore into the skin and advance at ten centimeters per hour – pretty fast for creatures less than one millimeter long! That’s why medical experts respectfully refer to them as “racing larvae.” Skin irritations usually occur in the vicinity of the worms.


They then bore through the wall of a blood vessel and are flushed into the lungs with the blood. Here, they penetrate the tissue and migrate upward through the trachea. “First coughed up, then swallowed – that’s how they enter the digestive tract,” explains Streit.

In the small intestine’s mucous membrane, each female lays up to a thousand unfertilized eggs per day. The majority are excreted in the feces. However, this human parasite has the unpleasant characteristic that a percent-



**Top** A free-living *Strongyloides papillosus* female, about one millimeter long: The mouth opening with the esophagus is at the top end. The intestine is connected to this. The animal lays its eggs through the vulva, located at the right center of the image.

**Bottom** The copulation organ of a free-living male *Strongyloides papillosus*. The small rod-shaped structures are bacteria clinging to it.



age of the embryos develop into infective larvae within the host. These penetrate the intestinal wall or bore through the anal mucous membrane and reenter the body. This is why, if left untreated, the infection can persist for a very long time.

In Cambodia, with the consent of the villagers, Streit's doctoral students Tegegn Jaleta and Siyu Zhou took fecal samples from humans and animals. "This was a huge event for the people

there," says Streit, smiling. "Many came to help." People were treated free of charge by staff from both institutes, and it was explained how they could protect themselves from infection.

The samples collected were first incubated for two days and placed in water, after which the worm larvae swimming in the water were separated out. However, genetic analyses weren't possible in the poorly equipped field laboratory in the village. The scientists thus had to fly the worms out to Germany, placing each one into a small tube containing ethanol. "Customs wasn't overly enthusiastic, but the officials were reassured once it became clear that the worms weren't being shipped alive and were in sterile packaging," explains Streit.

Streit and his colleagues then examined the worms' genetic material in their laboratory in Tübingen. The analysis revealed that one of the two *Strongyloides* populations that the researchers had identified in the dog feces was genetically identical to that in their owners' excrement. In other words, the

populations overlap, so dogs must be seriously considered as carriers.

As a next step, Streit wants to investigate whether dogs are the only carriers for humans. Water buffalo may also be candidates, as peasants in many areas still plough their rice fields with the animals, and do so barefoot. Water buffalo are already known to be carriers of other zoonotic diseases; studies in southern China, for example, revealed that the animals were the main carriers of schistosomiasis, a disease caused by trematodes.

"Although treatment of the patients was quite successful, the worms could hardly be suppressed by treating humans alone. Nevertheless, the infection was brought under control in China, but only once the water buffalo were also dewormed," says Streit.

Could this be a model for dealing with strongyloidiasis? To find this out, Streit is planning a project similar to the one in Cambodia, but this time in southwestern China. Here, not only are there rural areas where the threadworms are abundant, but there are also





**Top left** For their field study in Cambodia, the researchers from Tübingen collect stool samples from villagers, mix the samples with sawdust and cultivate them in Petri dishes. This allows the worms in the samples to continue their development.

**Top right** In a field laboratory, Siyu Zhou and Tegegn Jaleta prepare the samples for transportation to Germany, where the worms can be genetically analyzed.

**Bottom** Village life in northern Cambodia: People and animals live in close quarters and constantly come into contact with each other's excretions, allowing intestinal parasites to be easily transmitted.





» A warm, humid climate and a lack of hygiene are an El Dorado for worms.

highly qualified scientists with well-equipped laboratories.

A worm infection can be identified under the microscope. With more than 25,000 species worldwide, nematodes are hardly distinguishable for the layperson, but not for biologists. “Apart from *Parastrongyloides*, a close relative, *Strongyloides* is the only nematode in which the infective larvae have a very long esophagus typical of these species.

However, the fecal samples contain predominantly other nematodes, such as hookworms, as both humans and animals in Asia are often infected with various worm parasites. It remains unclear whether the different parasite species compete in the intestines. “But one thing is certain: in a further infection, worms can suppress new arrivals of the same species,” explains Streit. “We don’t yet know how, but this could offer an approach for future treatment.”

## RELEVANCE TO VETERINARY MEDICINE

To discover the worms’ tricks, Streit keeps two other *Strongyloides* species in Tübingen, where they live in rats or sheep (or, in the lab, in rabbits). Together with the University of Hohenheim, which maintains an animal breeding station in the Swabian Alb, he can also analyze the natural sheep parasite population for comparison.

In contrast to parasitic nematodes, *Strongyloides* doesn’t play a major role in veterinary medicine. Unlike *Strongyloides stercoralis*, other *Strongyloides*

species don’t lead to long-lasting, self-sustaining infections. However, these species, which can also live outside the host, are suitable as study objects for basic biological research.

Streit also wants to investigate whether the threadworms possess something like “parasite genes” – that is, a group of genes that are necessary for this lifestyle. In 2016, scientists decrypted the genomes of four different *Strongyloides* and two other closely related species, one of which occasionally lives parasitically, while the other is free-living. A genome comparison revealed that the parasitic worms have more genes for

two protein families that suppress the host’s immune response than their free-living relatives do. “We don’t yet know what role these genes play in a parasitic lifestyle,” Streit emphasizes. “For this, we would need to switch off the individual genes – not very easy in an organism that lives in a host every second generation.”

An additional peculiarity of these worms is that the free-living, bisexual generation produces exclusively female offspring, while the unisexual parasitic generation produces both males and females through parthenogenesis. Depending on the *Strongyloides* species,

## PARASITES AND THE IMMUNE SYSTEM

Many parasites suppress their hosts’ immune system, thus hindering a defense response. Long-standing host-parasite relationships can lead to an evolutionary race. This also applies to humans, who activate their immune system as a precaution to compensate for the suppressing effect of intestinal parasites.

This can become a problem if the immune system is constantly insufficiently challenged, for example in countries with high standards of hygiene. Scientists suspect that the immune system may then turn against its own body. This may explain why autoimmune diseases and allergies continue to increase in industrialized nations but are barely a problem in regions with many worm diseases. This is also suggested by the fact that parts of the immune system normally involved in parasite defense are overactive in autoimmune diseases and allergies.

This knowledge could be exploited medically. Some worm species are already being used to treat such autoimmune diseases as rheumatism, asthma, multiple sclerosis and Crohn’s disease, but there have not yet been any large-scale studies.



**Left** Adrian Streit spends most of his time researching in his laboratory in Tübingen. But in order to familiarize himself with the environmental conditions under which the threadworms live, he also conducts field studies – a welcome change from the daily routine at the institute.

**Right** Global distribution of *Strongyloides stercoralis*. The map shows that infection rates vary widely from country to country: more than 70 percent of the population may be infected with the worm in certain regions of some countries.

» Thanks to the two life cycles, a single, self-reproducing parasitic individual can establish a new population without sacrificing the benefits of sexual reproduction.

one chromosome must be completely or partially degraded so that males can develop without a father. The controlled degradation of genetic information is called *chromatin diminution* and was first identified in the equine roundworm. Such degradation is a rare occurrence in nature – outside of nematodes, it occurs, for example, in copepods, ciliates and lampreys.

### GENERATION WITHOUT MALES


Different *Strongyloides* species also appear to have different means of preventing males from developing in the progeny of the bisexual, free-living generation. Similar to many nematodes, the females of the rat parasite also have two X chromosomes, whereas the males have just one. A Y chromosome as in humans doesn't exist in these species.

“While we didn't find mature sperm leading to males in sheep worms, there are sperm with and without an X chromosome in rat parasites,” Streit explains. Some of these sperm would therefore have to lead to males, since an egg cell always carries an X chromosome. In contrast to the sheep, there are actually male worm embryos in the rat. However, they apparently die off, because neither of the two worm species forms male larvae. Still, when the male-designated sperm or the early male embryos are screened out remains a mystery.

Why is a parasite's life so complicated? As Adrian Streit sees it, the complex life cycle emerged gradually from a simpler one: numerous free-living nematodes occasionally form permanent stages to survive hard times. They often attach themselves to other ani-

mals, such as the previously mentioned worm living on a beetle. “Once such a larva is attached to an animal, the step to get inside the animal is no longer that great. This may lead to a life cycle similar to that of *Strongyloides*, where the worm is either parasitic or free-living,” explains Streit. However, in the majority of parasitic nematodes, every generation is parasitic. Thus, in the course of evolution, the free-living phase may have been sacrificed in favor of a purely parasitic one in many parasites.

*Strongyloides* may be following the same path. But has it possibly taken a wrong turn? After all, the females multiply in their hosts exclusively by unisexual parthenogenesis, so “rejuvenation” of the genome by gene recombination doesn't take place in the parasitic generation.



In evolutionary terms, unisexually or asexually reproducing lines are generally young. It may even be assumed that they can't age, because the transition to a life without sexual reproduction is the beginning of the end. In other words, *Strongyloides* may have maneuvered itself into a cul-de-sac, making it impossible for it to now rescind the sexual free-living cycle.

But maybe the worm has found an ideal solution for itself: thanks to the two life cycles, a single, self-reproducing parasitic individual can establish a new population without sacrificing the benefits of sexual reproduction.

Whether and how *Strongyloides* will ever become a pure parasite, or whether its life cycle will become even more complicated, can't be predicted with certainty today. Evolution always finds a new, and sometimes curious, solution. ◀

📍 [www.tinyurl.com/yczguvf9](http://www.tinyurl.com/yczguvf9) (available only in German)

### TO THE POINT

- There are more than 50 species of parasitic *Strongyloides* (a threadworm) that infect a wide range of terrestrial vertebrates. An estimated 300 million people worldwide are infected with the *Strongyloides stercoralis* worm.
- Dogs can also be infected with *Strongyloides stercoralis*. The parasites can infect people through the animals' feces. The infection is usually harmless in humans, but it can end in death in individuals with a weakened immune system.
- Genetic analyses have revealed that parasitic worms have more genes compared with free-living species, potentially reducing the host's immune response.

### GLOSSARY

**Parthenogenesis:** This type of unisexual reproduction leads to progeny from unfertilized eggs produced by female animals. In mitotic parthenogenesis, such as occurs in *Strongyloides*, diploid oocytes are formed without meiosis, and the genome isn't remixed by recombination. In other words, it is a form of asexual reproduction and the progeny are therefore genetically identical to their mother.